Tanacetum cinerariifolium

Habitus and growth type

Plant height [m]: **0.5**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Temperature indicator value: **9**

Disturbance indicator values

Disturbance frequency: **0.64**
Disturbance frequency (herb layer): **1.62**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.2**
Grazing pressure: **0.4**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz