Tanacetum corymbosum

Habitus and growth type

Plant height [m]: **0.73**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.83**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.47**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**
Temperature indicator value: **6.3**
Reaction indicator value: **7.5**
Nutrient indicator value: **3.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.14**
Disturbance frequency (herb layer): **1.38**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.2**
Mowing frequency: **0.1**
Grazing pressure: **0.22**
Soil disturbance: **0.11**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**, **T19 Temperate and submediterranean thermophilous deciduous forest**
Constant species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**, **T19 Temperate and submediterranean thermophilous deciduous forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz