Tanacetum vulgare

Habitus and growth type

Plant height [m]: **0.81**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **20.61**

Flower

Flowering period: **June-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.21**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Reaction indicator value: **7.1**
Nutrient indicator value: **5.7**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.15**
Disturbance frequency (herb layer): **1.63**
Disturbance severity: **0.62**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.58**
Mowing frequency: **0.7**
Grazing pressure: **0.17**
Soil disturbance: **0.3**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **V38 Dry perennial anthropogenic herbaceous vegetation**
Constant species of EUNIS habitats: **V38 Dry perennial anthropogenic herbaceous vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz