# *Teucrium divaricatum*

## Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

## Fruit, seed and dispersal

Seed mass [mg]: **1.91**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **11**
Moisture indicator value: **2**
Reaction indicator value: **8**
Salinity indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **0.16**
Disturbance frequency (herb layer): **1.48**
Disturbance severity: **0.72**
Disturbance severity (herb layer): **0.23**
Mowing frequency: **0.02**
Grazing pressure: **0.3**
Soil disturbance: **0.15**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**

## Broad habitat

Occurrence in broad habitats: **Scrub**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*