Teucrium microphyllum

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.26**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**
Moisture indicator value: **2**
Reaction indicator value: **7**
Nutrient indicator value: **2**
Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.5**
Disturbance frequency (herb layer): **1.77**
Disturbance severity: **0.74**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.09**
Grazing pressure: **0.38**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**, **S72 Eastern Mediterranean spiny heath (phrygana)**

Constant species of EUNIS habitats: **S63 Eastern garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz