Trifolium alpestre

Habitus and growth type

Plant height [m]: **0.26**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **11.28**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.62**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Temperature indicator value: **5.8**
Moisture indicator value: **3.4**
Reaction indicator value: **6.6**
Nutrient indicator value: **3.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.44**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.53**
Disturbance severity: **0.55**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.33**
Grazing pressure: **0.25**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

EUNIS habitat

Constant species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**, **R51 Thermophilous forest fringe of base-rich soils**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz