# *Trifolium strictum*

## Habitus and growth type

Plant height [m]: **0.1**
Life span: **Annual**
Life form: **Therophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.45**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Oligotrophic**

### Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **7.3**
Moisture indicator value: **2**
Nutrient indicator value: **1.5**
Salinity indicator value: **2.3**

### Disturbance indicator values

Disturbance frequency: **1.67**
Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.66**
Grazing pressure: **0.34**
Soil disturbance: **0.25**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea*, FH (BUL) *Poetea bulbosae*, FI (TUB) *Helianthemetea guttati***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R24 Iberian summer pasture (vallicar), S34 Balkan-Anatolian submontane genistoid scrub**

Constant species of EUNIS habitats: **R24 Iberian summer pasture (vallicar), S34 Balkan-Anatolian submontane genistoid scrub**

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Saline vegetation, Scrub**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*