

Veratrum lobelianum

Habitus and growth type

Plant height [m]: **0.92**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.74**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **3.3**

Moisture indicator value: **6.2**

Reaction indicator value: **6**

Nutrient indicator value: **6.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**

Disturbance frequency (herb layer): **1.12**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.09**

Grazing pressure: **0.19**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CS \(MUL\) *Mulgedio-Aconitetea***](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**Q23 Relict mire of Mediterranean mountains**](#), [**R56 Montane to subalpine moist or wet tall-herb and fern fringe**](#), [**S25 Subalpine and subarctic deciduous scrub**](#)

Constant species of EUNIS habitats: [**Q23 Relict mire of Mediterranean mountains**](#), [**R56 Montane to subalpine moist or wet tall-herb and fern fringe**](#), [**S25 Subalpine and subarctic deciduous scrub**](#)

Broad habitat

Occurrence in broad habitats: **Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub**