# *Veronica anagalloides*

### Habitus and growth type

Plant height [m]: **0.27**
Life span: **Annual**
Life form: **Therophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **19.58**

### Flower

Flowering period: **May-October**

### Fruit, seed and dispersal

Seed mass [mg]: **0.03**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **7.7**
Moisture indicator value: **8.7**
Salinity indicator value: **0.5**

#### Disturbance indicator values

Disturbance frequency: **1.7**
Disturbance frequency (herb layer): **1.92**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.57**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.2**
Grazing pressure: **0.23**
Soil disturbance: **0.27**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

### Broad habitat

Occurrence in broad habitats: **Spring, Wetland**

## Distribution

Continentality: **7**
Continentality amplitude: **6**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*