Veronica gentianoides

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **12.92**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **5.3**
Moisture indicator value: **6.5**

Disturbance indicator values

Disturbance frequency: **1.49**
Disturbance frequency (herb layer): **1.72**
Disturbance severity: **0.34**
Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.11**
Grazing pressure: **0.26**
Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of phytosociological classes: **AB (KOB) *Carici rupestris-Kobresietea bellardii***

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz