Bupleurum commutatum

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.9**
Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7.8**
Moisture indicator value: **3.1**
Reaction indicator value: **6.5**
Nutrient indicator value: **3.9**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **2.23**
Disturbance frequency (herb layer): **2.23**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.49**
Grazing pressure: **0.37**
Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz