Beta vulgaris subsp. *maritima*

Fruit, seed and dispersal

Seed mass [mg]: **16.44**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Ecology

Disturbance indicator values

Disturbance frequency: **1.54**
Disturbance frequency (herb layer): **1.61**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.63**
Mowing frequency: **0.23**
Grazing pressure: **0.18**
Soil disturbance: **0.54**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JC (CAK) *Cakiletea maritimae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N21 Atlantic, Baltic and Arctic coastal shingle beach**, **N22 Mediterranean and Black Sea coastal shingle beach**, **N31 Atlantic and Baltic rocky sea cliff and shore**
Constant species of EUNIS habitats: **N21 Atlantic, Baltic and Arctic coastal shingle beach**, **N22 Mediterranean and Black Sea coastal shingle beach**, **N31 Atlantic and Baltic rocky sea cliff and shore**
Dominant species of EUNIS habitats: **N21 Atlantic, Baltic and Arctic coastal shingle beach**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Coastal cliff**

Distribution

Continentality: **3**
Continentality amplitude: **3**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz