Campanula latifolia subsp. *latifolia*

Habitus and growth type

Plant height [m]: **0.83**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **55.58**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.8**
Temperature indicator value: **4.8**
Moisture indicator value: **6.1**
Reaction indicator value: **7.7**
Nutrient indicator value: **6.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.12**
Disturbance frequency (herb layer): **1.35**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.03**
Grazing pressure: **0.18**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae*, CS (MUL) *Mulgedio-Aconitetea***

Broad habitat

Occurrence in broad habitats: **Wetland, Alpine-subalpine and arctic grassland, Scrub, Forest, Synanthropic**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz