Festuca stricta subsp. *sulcata*

Habitus and growth type

Plant height [m]: **0.39**

Fruit, seed and dispersal

Seed mass [mg]: **0.46**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Ecology

Disturbance indicator values

Disturbance frequency: **1.11**
Disturbance frequency (herb layer): **1.82**
Disturbance severity: **0.5**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.74**
Grazing pressure: **0.28**
Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei*, DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1A Semi-dry perennial calcareous grassland (meadow steppe), R1B Continental dry grassland (true steppe), R1C Desert steppe, S36 Low steppic scrub**
Constant species of EUNIS habitats: **R16 Perennial rocky grassland of Central and South-Eastern Europe, R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region, R1A Semi-dry perennial calcareous grassland (meadow steppe), R1B Continental dry grassland (true steppe), R1C Desert steppe, S36 Low steppic scrub, T3H Larix light taiga**
Dominant species of EUNIS habitats: **R1A Semi-dry perennial calcareous grassland (meadow steppe)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **8**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz