

Festuca stricta subsp. *trachyphylla*

Habitus and growth type

Plant height [m]: **0.26**

Leaf

Specific leaf area [mm²/mg]: **5.65**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Ecology

Disturbance indicator values

Disturbance frequency: **1.27**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.74**

Grazing pressure: **0.31**

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) *Koelerio-Corynephoretea canescentis*](#), [DA \(FES\) *Festuco-Brometea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils](#)

Constant species of EUNIS habitats: [R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils](#)

Dominant species of EUNIS habitats: [R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **2**