Festuca stricta subsp. *trachyphylla*

Habitus and growth type

Plant height [m]: **0.26**

Leaf

Specific leaf area [mm^2/mg]: **5.65**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **1, 7**

Ecology

Disturbance indicator values

Disturbance frequency: **1.27**
Disturbance frequency (herb layer): **1.92**
Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.74**
Grazing pressure: **0.31**
Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis*, DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**
Constant species of EUNIS habitats: **R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**
Dominant species of EUNIS habitats: **R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz