Chaenorhinum litorale subsp. *pterosporum*

Habitus and growth type

Plant height [m]: **0.4**
Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Temperature indicator value: **8**

Disturbance indicator values

Disturbance frequency: **1.39**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.68**
Mowing frequency: **0.57**
Grazing pressure: **0.15**
Soil disturbance: **0.43**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea***, **KE (THL) *Thlaspietea rotundifolii***

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz