Lysimachia nemorum subsp. *nemorum*

Habitus and growth type

Plant height [m]: **0.27**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **30.2**

Fruit, seed and dispersal

Seed mass [mg]: **0.49**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **7.7**
Reaction indicator value: **5.7**
Nutrient indicator value: **5.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.2**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.09**
Grazing pressure: **0.19**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1T Azorean open, dry, acid to neutral grassland**, **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**

Constant species of EUNIS habitats: **R1T Azorean open, dry, acid to neutral grassland**, **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **3**

Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz