Panicum capillare subsp. *capillare*

Habitus and growth type

Plant height [m]: **0.34**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **28.77**

Fruit, seed and dispersal

Seed mass [mg]: **0.36**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**
Moisture indicator value: **4**
Reaction indicator value: **6.8**
Nutrient indicator value: **5.4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.63**
Disturbance frequency (herb layer): **1.77**
Disturbance severity: **0.8**
Disturbance severity (herb layer): **0.8**
Mowing frequency: **0.34**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.18**
Soil disturbance: **0.48**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QD (DIG) *Digitario sanguinalis-Eragrostietea minoris***

Broad habitat

Occurrence in broad habitats: **Wetland, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz