Salix cinerea subsp. *cinerea*

Habitus and growth type

Plant height [m]: **4.7**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **0.17**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Ecology

Disturbance indicator values

Disturbance frequency: **0.54**
Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.13**
Grazing pressure: **0.18**
Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S92 Salix fen scrub**, **T15 Broadleaved swamp forest on non-acid peat**, **T16 Broadleaved mire forest on acid peat**
Constant species of EUNIS habitats: **S92 Salix fen scrub**, **T15 Broadleaved swamp forest on non-acid peat**, **T16 Broadleaved mire forest on acid peat**
Dominant species of EUNIS habitats: **S92 Salix fen scrub**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz