Salix cinerea subsp. *oleifolia*

Habitus and growth type

Plant height [m]: **4.7**

Leaf

Specific leaf area [mm^2/mg]: **12.3**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **0.17**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Ecology

Disturbance indicator values

Disturbance frequency: **0.53**
Disturbance frequency (herb layer): **1.47**
Disturbance severity: **0.55**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.09**
Grazing pressure: **0.22**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FD (NER) *Nerio-Tamaricetea*, HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S91 Temperate riparian scrub, T14 Mediterranean and Macaronesian riparian forest**
Constant species of EUNIS habitats: **S91 Temperate riparian scrub, T14 Mediterranean and Macaronesian riparian forest**
Dominant species of EUNIS habitats: **S91 Temperate riparian scrub**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz