Schoenoplectus lacustris subsp. *glaucus*

Habitus and growth type

Plant height [m]: **1.18**

Leaf

Specific leaf area [mm^2/mg]: **5.42**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.54**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Ecology

Disturbance indicator values

Disturbance frequency: **1.83**
Disturbance frequency (herb layer): **1.9**
Disturbance severity: **0.34**
Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.45**
Grazing pressure: **0.25**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q46 Carpathian travertine fen with halophytes**, **Q54 Inland saline or brackish helophyte bed**
Constant species of EUNIS habitats: **MA232 Baltic coastal meadow**, **Q46 Carpathian travertine fen with halophytes**, **Q54 Inland saline or brackish helophyte bed**
Dominant species of EUNIS habitats: **Q46 Carpathian travertine fen with halophytes**, **Q54 Inland saline or brackish helophyte bed**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Wetland, Mire, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution

Continentality: **7**
Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz